

Access this article online
Quick Response Code:

Website: www.jehp.net
DOI: 10.4103/jehp.jehp_41_17

Estimating the carbohydrate content of various forms of tobacco by phenol-sulfuric acid method

Vardhaman Mulchand Jain, Gundabaktha Nagappa Karibasappa¹, Arun Suresh Dodamani, Gaurao Vasant Mali

Abstract:

BACKGROUND: Due to consumption of various forms of tobacco in large amounts by Indian population, it has become a cause of concern for major oral diseases. In 2008, the WHO named tobacco as the world's single greatest cause of preventable death. It is also known that certain amount of carbohydrates are incorporated in processed tobacco to make it acceptable for consumption. Thus, its role in oral diseases becomes an important question at this point of time. Through this study, it is attempted to find out the carbohydrate content of various forms of tobacco by phenol-sulfuric acid method.

MATERIALS AND METHODS: Tobacco products selected for the study were Nandi hookah tambakhu (A), photo brand budhaa Punjabi snuff (B), Miraj (C), Gai-chhap tambakhu (D), Hanuman-chhap Pandharpuri tambakhu (E), and Hathi-chhap Bidi (F). The samples were decoded and transported to laboratory and tested at various concentrations by phenol-sulfuric acid method followed by ultraviolet spectrophotometry to determine their absorbance.

RESULTS: The present study showed Hathi-chhap bidi/sample F had a maximum absorbance (1.995) at 10 µg/ml which is a smoking form of tobacco followed by rest all smokeless forms of tobacco, i.e. sample C (0.452), sample B (0.253), sample D (0.077), sample E (-0.018), and sample A (-0.127), respectively.

CONCLUSION: As the concentration of tobacco sample increases, their absorbance increases which in turn is suggestive of increase in its carbohydrate concentration. Carbohydrates in the form of sugars, either inherently present or added in it during manufacturing can serve as a risk factor for higher incidence of dental caries.

Keywords:

Carbohydrate, caries, phenol-sulfuric acid, spectrophotometry, tobacco

Department of Public Health Dentistry, ACPM Dental College, Dhule,
¹Department of Public Health Dentistry, Ajeenkya DY Patil Dental School, Pune, Maharashtra, India

Address for correspondence:

Dr. Vardhaman Mulchand Jain, Assistant Professor, Department of Public Health Dentistry, ACPM Dental College, P. Box No. 145, Opposite Jawahar Soot Girmi, Sakri Road, Dhule - 424 001, Maharashtra, India.
E-mail: dr.vardhaman.jain@gmail.com

Received: 25-04-2017
Accepted: 13-06-2017

Introduction

India is the second largest consumer of tobacco in the world and majority of Indian population consume tobacco in various forms.^[1] Tobacco is the most easily accessible, legally available addictive substance which contributes significantly to premature death and long-term suffering, and also being a major risk factor for cardiovascular diseases, chronic obstructive pulmonary

diseases, cancers, reproductive outcomes, and oral diseases.^[2]

In India, tobacco is consumed in myriad forms which include smoking as well as smokeless tobacco (SLT).^[3] Bidi is the most popular prevalent smoking product consumed in rural areas in comparison to cigarette smoking more preferably used in urban areas. Hookah, chuttas, dhumti, chillum, cigars, cheroots, and pipes are some other smoking forms of tobacco in different parts of India.^[2]

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Jain VM, Karibasappa GN, Dodamani AS, Mali GV. Estimating the carbohydrate content of various forms of tobacco by phenol-sulfuric acid method. *J Edu Health Promot* 2017;6:90.

SLT is consumed predominantly by chewing in form of pan (piper betel leaf filled with sliced areca nut, lime, catechu, and other spices chewed with or without tobacco), pan-masala or gutkha (a chewable tobacco containing arecanut), and mawa in many parts of India. Furthermore, tobacco such as mishri (a powdered tobacco rubbed on the gums as toothpaste), gul, and gudakhu are widely used as topical application on teeth and gums.^[3,4]

The prevalence of various forms of tobacco usage and habit varies across different geographical areas. It has been predicted that if the current trends of tobacco use are unchecked the number of tobacco-related deaths will reach eight million by 2030, with approximately 70% of these deaths occurring in developing countries. SLT use affects nearly 300 million people worldwide and about 90% of the world's users reside in the South-East Asian Region of the WHO.^[5]

The term "additive" means any substance, intended for use as a flavoring or coloring or in producing, manufacturing, packing, processing, preparing, treating, packaging, transporting, or holding, etc.^[6]

Carbohydrates are the natural tobacco components present in levels up to 20 wt% in tobacco products and used as additives during manufacturing process. They also serve as a flavoring substance and humectant in tobacco products.^[7]

Due to the consumption of tobacco in such large amounts, it has become a cause of concern for major oral diseases. There is abundance literature available depicting the role of carbohydrates in causation and progression of dental caries.^[8,9] Tobacco addicts in any form are prone to dental caries process.^[10] It might be due to continuous consumption of these products multiple times and for chronic duration. Many studies have assessed the carbohydrate content of smoked forms of tobacco; however, there is scarcity of evidence that assesses the carbohydrate content of smokeless forms of tobacco.^[7]

Phenol-sulfuric acid method is the one of the most easiest and reliable, colorimetric method, widely used among the quantitative assays available for carbohydrate estimation in aqueous solutions.^[11-13] It is followed by determining light absorbance on ultraviolet (UV) spectrophotometer which is one of the oldest techniques of analysis and is the basis for number of ideal methods for the determination of micro and semi-micro quantity of analytes in a sample.^[14]

The study area Khandesh being a low socioeconomic region, bidi is the most common smoked form of tobacco

consumed along with numerous other smokeless forms of tobacco.^[15] Hence, the present study was conducted to assess and compare the carbohydrate content of different tobacco products in smoked and smokeless forms by phenol-sulfuric acid method.

Materials and Methods

The present study is an *in vitro* study conducted at H. R. Patel Institute of Pharmaceutical Education and Research at Shirpur, Maharashtra. Ethical clearance for the study was obtained from the Institutional Ethical Review Committee.

Selection and coding of sample

The various commercially available tobacco products were taken into consideration based on their popularity, widespread use among the people, availability in the vicinity of study (Dhule) area, and of those, the following samples were randomly selected which were labeled in the following manner:

- A. Nandi hookah tambakhu
- B. Photo brand budhaa Punjabi snuff
- C. Miraj
- D. Gai-chhap tambakhu
- E. Hanuman-chhap Pandharpuri tambakhu
- F. Hathi-chhap Bidi.

Recently, manufactured and packed samples were purchased from the local retail outlet. The study was conducted over a period of 15 days.

Packing and transport

The samples were weighed according to 100 mg of tobacco content and were sealed in air-tight pouches with labeling of the codes (A, B, C, D, E, F). The samples were then transported in hygienic and moisture-proof conditions to the study center.

Processing

Phenol-sulfuric acid method is the most reliable method among all the quantitative assays for carbohydrate estimation.^[11-13] In hot acidic medium, glucose is dehydrated to hydroxymethyl furfural. This forms a yellow-brown-colored product with phenol and has absorption maximum at 490 nm UV spectroscopy.^[12,16] This obeys Beer-Lambert's law which states that when a beam of monochromatic light is passed through a solution of an absorbing substance, the rate of decrease of intensity of radiation with thickness of absorbing solution is proportional to incident radiation as well as the concentration of solution.^[17]

- $A = \log(I_0/I) = \epsilon cL$
- I_0 = Intensity of light incident upon sample
- I = Intensity of light leaving sample cell
- c = Molar concentration of solute

- L = Length of sample cell (cm)
- E = Molar absorptivity
- A = Absorbance.

Apparatus

The apparatus used in the study were pipettes (micropipette), UV spectrophotometer (Shimadzu, Asia Pacific Pvt. Ltd., Singapore), reflux, condenser, test tube, stand, boiler, cuvette, etc.

Reagents

96% sulfuric acid, standard glucose, 2.5 N HCl, sodium carbonate, distilled H₂O, phenol solution.

Procedure for standardized sample

Take 100 mg of glucose into test tube. Add 5 ml of 2.5 N HCl and boil in water bath for 3 h to hydrolyze the sugars. Cool it to room temperature. Add sufficient quantity of solid sodium carbonate until the effervescence ceases. This indicates complete neutralization. Filter and make the volume up to 100 ml. Pipette out 0.2, 0.4, 0.6, 0.8, and 1 ml of working standard into a series of test tubes.

The blank was set initially with all reagents without sample. Add 1 ml of phenol solution to each tube and shake well. Then, add 5 ml of sulfuric acid 96% and again shake well. Keep aside for 10 min. Shake each test tube and keep in water bath at 25°C–30°C for 20 min. Now measure the color at 490 nm. After that switch on the spectrometer, first take the absorbance (optical density [OD]) of blank and make it zero. Take the OD of all the test tubes and wash the cuvette each time after taking OD.

The values obtained of absorbance were helpful to plot a graph which is very crucial in estimating carbohydrate content. The values should be positive, above zero to plot a graph. After the plotting of a graph, the total amount of carbohydrate in the sample from glucose standard graph was calculated. The standard curve of absorbance was plotted at 490 nm on “Y” axis representing absorbance at 490 nm versus concentration of glucose in µg/ml on X axis [Figure 1].

From the above graph, we get an equation,

$$Y = mx + c$$

- Y = Absorbance of drug at 490 nm
- m = Mass
- x = Carbohydrate concentration of drug
- c = Velocity constant.

$$Y = 0.195x + 0.029$$

$$(Y - 0.029)/0.195 = x$$

- Absorbance corresponds to 0.1 ml of the test = x mg of glucose

- One hundred milliliter of the sample solution contains = $x/0.1 \times 100$ mg of glucose = % of total carbohydrate present in it.^[11,18]

Procedure for tobacco samples to be tested:

To have standard comparison of each sample, the procedure carried out for glucose sample was repeated in similar fashion with all the tobacco samples and their absorbance was determined at concentrations of 0.1, 2.5, 5, 10 µg/ml.

Results

The present study shows that Hathi-chhap bidi/sample F has maximum absorbance (1.995) at 10 µg/ml which is a smoking form of tobacco followed by rest all smokeless forms of tobacco, i.e. sample C (0.452), sample B (0.253), sample D (0.077), sample E (-0.018), and sample A (-0.127), respectively [Table 1].

Similarly, sample F showed maximum absorbance followed by sample C, sample B, sample D, sample E, and sample A, respectively, at different concentration of 0.1, 2.5, 5 µg/ml [Table 1].

Discussion

In the present study, maximum absorption was seen with bidi followed by other smokeless forms of tobacco,




Figure 1: Standard curve of absorbance at 490 nm on “Y” axis representing absorbance at 490 nm versus concentration of glucose in µg/ml on X axis. From the above graph, we get an equation, $Y = mx + c$. Y = Absorbance of drug at 490 nm, m = Mass, x = Carbohydrate concentration of drug, c = Velocity constant

Table 1: Absorbance of tobacco samples at different concentrations

Concentration (µg/ml)	Absorbance					
	A	B	C	D	E	F
0.1	-1.995	-0.095	-0.058	-0.362	-0.428	-0.032
2.5	-0.427	-0.060	-0.053	-0.108	-0.236	-0.014
5	-0.192	0.005	0.146	-0.018	-0.066	0.534
10	-0.127	0.253	0.452	0.077	-0.018	1.995

i.e., sample C, B, D, E, and A, respectively. This in turn suggests that smoking form of tobacco has more carbohydrate content compared to smokeless forms of tobacco.

Phenol-sulfuric acid method was used for determining carbohydrate concentration as it is one of the most widely used colorimetric methods till date.^[11] The basic principle of this method is that carbohydrates, when dehydrated by reaction with concentrated sulfuric acid, produce furfural derivatives, which further reacts with phenol to develop detectable color. Then, light absorption at 490 nm is recorded on a spectrophotometer.^[12,16]

Carbohydrates are polyhydroxy compounds consisting of aldose or ketose sugar. Hence, placing the tobacco products for longer time in oral cavity increases the risk of dental caries, i.e. sugar and oral health are integrally related to each other.^[19] Furthermore, more the frequency of tobacco consumption, more will be the risk for sugar exposure to that individual. This provided an impetus to estimate the carbohydrate in various tobacco products available in the near vicinity of study area.

We could not plot a standard absorbance curve at 490 nm of the tested samples as some of the values were negative, so only descriptive statistics for absorbance is given at different concentrations. As we could not plot the graph, $Y = mx + c$ equation was not obtained, and hence, the carbohydrate content readings were not given by spectrophotometer. As the absorbance is directly proportional to carbohydrate content, it is predicted in the present study that as the absorbance increases, the carbohydrate content increases.

Among the samples tested, bidi which is smoking form of tobacco had more carbohydrate content compared to other smokeless forms of tobacco. It is supported by the findings of Jansen *et al.*^[7] and Ramusino *et al.*^[20]

Among the smokeless forms, sample C (Miraj) had maximum carbohydrate followed by sample B, D, E, and A, respectively. We could not compare the present findings with other similar studies as there were no such studies available in the previous literature. Many studies in the previous literature were conducted on smoking form of tobacco and there is paucity of literature regarding smokeless forms of tobacco.

The literature associating SLT use with either increasing or decreasing dental caries incidence is even harder to find compared to the literature associating smoking form of tobacco with dental caries. Theories have been postulated based on limited clinical findings, chemical analysis of the content of various ST products, and

in vitro effects of ST on the growth of bacteria that has been implicated in caries development.^[21]

Evidence linking ST use with increased dental caries prevalence has been reported.^[22,23] In a case report by Croft, a 54-year-old patient presented “cervical caries” in the area of tobacco placement and gingivitis and recession with same tooth.^[23,24] In contrast, Zitterbart *et al.* did not find any evidence of caries in the area of quid placement in their 36-year-old tobacco chewer.^[23,25] Another study, which was performed among Swedish children did not report any prevalence of caries among snuff users.^[26] On the other hand, higher prevalence of caries was observed in snuff dippers than in nontobacco users among teenagers in Gothenburg.^[27]

There is abundance of literature which states that sugar is responsible for causing dental caries,^[21,28-30] but in the present study, very less amount of sugar is present in tobacco products which might be inherently present or added in it. This shows that sugar is not responsible for directly causing caries in these individuals. Even though ST do not directly cause dental caries, placing tobacco product for longer duration in the oral cavity may lead to chronic irritation which in turn causes gingival recession, denudation of root surface of teeth that might increase the risk for occurrence of root caries.^[10,31-33]

Although there is insufficient evidence, to conclude that SLT has a direct causal role in either caries formation or inhibition, literature suggests SLT do play an important role in caries activity and are harmful with respect to an individual’s health.

Review of studies conducted on oral consequences of snuff and chewing tobacco use among professional baseball players in US found that ST use showed a significantly higher prevalence of root caries compared to nonsmokers.^[34] Data from the multipurpose health survey conducted in USA showed that the mean number of decayed and filled root surface for those who used chewing tobacco was four times higher than for those who did not use tobacco. It is important to note that the decayed or filled surfaces tended to match the side of mouth on which the ST was used. The results showed that the mean number of decayed and filled root surface increased with increasing number of chewing tobacco packets used per week and duration of its use.^[35]

The results obtained in the present study show that the absorbance was low of all the smokeless forms of tobacco compared to smoking form of tobacco which indicates very low or negligible quantity of carbohydrate present in it comparatively. This suggests less amount of inherent sugars as well as additives present in smokeless forms compared to smoking

form of tobacco. The types of sweeteners and sugars commonly found in ST are fructose, glucose, sucrose, maltose, and isomaltose.^[36] This addition is presumed to be having a neutralizing effect on the bitter taste of tobacco.^[37] Large variations in sugar among tobacco products can exist within form-to-form, brand-to-brand, and state-to-state. This may explain the diverse opinions of dental practitioners and investigators with respect to the concept of tobacco, increasing or decreasing incidence of dental caries.

The present study had its own limitations such as exact carbohydrate content of the samples was not estimated as some values obtained were negative due to which graph could not be plotted. The present study was done by phenol-sulfuric acid method taking financial constraint into consideration. Further research has to be conducted to determine carbohydrate content of different forms of SLT by various other techniques such as chromatography, capillary electrophoresis, infrared spectroscopy, light scattering detection, and nuclear magnetic resonance spectroscopy.

Conclusion

As the concentration of tobacco increases, the absorbance of the tobacco products (sample) increases which in turn is suggestive of increase in its carbohydrate concentration.

- Among the tested samples, maximum absorbance was seen with sample F (Bidi) which is a smoking form of tobacco
- Bidi had more absorbance that indicates more carbohydrate content compared to smokeless forms of tobacco
- Among the smokeless forms of tobacco, sample C (Miraj) had maximum absorbance that indicates maximum carbohydrate content followed by sample B, D, E, and A, respectively
- Carbohydrates in the form of sugars, either inherently present or added in it during manufacturing can serve as contributing risk factor for higher incidence of dental caries.

Acknowledgment

We would like to thank all the staff members and lab technician, H. R. Patel Institute of Pharmaceutical Education and Research, Shirpur, Maharashtra, for helping them to successfully conduct the study.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

References

1. Mishra GA, Pimple SA, Shastri SS. An overview of the tobacco problem in India. *Indian J Med Paediatr Oncol* 2012;33:139-45.
2. Bhawna G. Burden of smoked and smokeless tobacco consumption in India – Results from the Global adult Tobacco Survey India (GATS-India)-2009-2010. *Asian Pac J Cancer Prev* 2013;14:3323-9.
3. Rani M, Bonu S, Jha P, Nguyen SN, Jamjoum L. Tobacco use in India: Prevalence and predictors of smoking and chewing in a national cross sectional household survey. *Tob Control* 2003;12:e4.
4. Gupta PC, Ray CS. Smokeless tobacco and health in India and South Asia. *Respirology* 2003;8:419-31.
5. Sinha DN, Rizwan SA, Aryal KK, Karki KB, Zaman MM, Gupta PC. Trends of smokeless tobacco use among adults (aged 15-49 years) in Bangladesh, India and Nepal. *Asian Pac J Cancer Prev* 2015;16:6561-8.
6. Family Smoking Prevention and Tobacco Control Act. [Internet]. [cited 2016 Jan 2]. Available from: <http://www.fda.gov/tobaccoproducts/labeling/.../ucm237092.htm>.
7. Jansen E, Cremers J, Borst S, Talhout R. Simple determination of sugars in cigarettes. *J Anal Bioanal Tech* 2014;5:1-3.
8. Nizel Abraham E, Papas Athena S. The Role of Carbohydrates in the Production of Dental Caries. *Nutrition in Clinical Dentistry*, 3rd ed. Philadelphia: W. B. Saunders Company; 1989.p. 31-50.
9. Mandel ID. Effects of dietary modifications on caries in humans. *J Dent Res* 1970;49:1201-11.
10. Going RE, Hsu SC, Pollack RL, Haugh LD. Sugar and fluoride content of various forms of tobacco. *J Am Dent Assoc* 1980;100:27-33.
11. Dubois M, Gilles K, Hamilton J, Rebers P, Smith F. Colorimetric method for determination of sugars and related substances. *Analyt Chem* 1956;28:350-6.
12. HiPer® Carbohydrates Estimation Teaching Kit (Quantitative). Available from: <http://www.himedialabs.com/TD/HTBC003.pdf>. [Last accessed on 2016 Jan 16].
13. Masuko T, Minami A, Iwasaki N, Majima T, Nishimura S, Lee YC. Carbohydrate analysis by a phenol-sulfuric acid method in microplate format. *Anal Biochem* 2005;339:69-72.
14. Introduction to UV-Visible Spectroscopy. Available from: <http://www.slideshare.net/JunaidKhan30/introduction-to-uv-visible-spectroscopy>. [Last accessed on 2016 Jan 16].
15. Pednekar MS, Gupta PC, Yeole BB, Hébert JR. Association of tobacco habits, including bidi smoking, with overall and site-specific cancer incidence: Results from the Mumbai cohort study. *Cancer Causes Control* 2011;22:859-68.
16. Sadasivam S, Manickam A. Phenol sulphuric acid method for total carbohydrate. In: *Biochemical Methods*. New Delhi: New Age International (P), Ltd.; 2005. p. 10.
17. Basic UV-Vis Theory, Concepts and Applications. Available from: <http://www.molecularinfo.com/MTM/UV.pdf>. [Last accessed on 2016 Jan 16].
18. Agrawal N, Divya K, Minj DK, Rani K. Estimation of total carbohydrate present in dry fruits. *IOSR-J Environ Sci Toxicol Food Technol* 2015;1:24-7.
19. Moynihan P, Petersen PE. Diet, nutrition and the prevention of dental diseases. *Public Health Nutr* 2004;7:201-26.
20. Ramusino MC, Dattilo BS, Lucibello A, Rossi SG. Determination of 25 low molecular weight carbohydrates in tobacco by high performance ion chromatography. *Beitrag Zur Tabakforschung International/Contributions to Tobacco research* 1994;16:77-84.
21. Vellappally S, Fiala Z, Smejkalová J, Jacob V, Shriharsha P. Influence of tobacco use in dental caries development. *Cent Eur J Public Health* 2007;15:116-21.

22. Sitzes L Jr. On chewing tobacco [letter]. *ADA News* 1977;8:6.
23. Weintraub JA, Burt BA. Periodontal effects and dental caries associated with smokeless tobacco use. *Public Health Rep* 1987;102:30-5.
24. Croft L. Smokeless tobacco. Case report II. *Tex Dent J* 1983;100:14-5.
25. Zitterbart PA, Marlin DC, Christen AG. Dental and oral effects observed in a long-term tobacco chewer: Case report. *J Indiana Dent Assoc* 1983;62:17-8.
26. Modéer T, Lavstedt S, Ahlund C. Relation between tobacco consumption and oral health in Swedish schoolchildren. *Acta Odontol Scand* 1980;38:223-7.
27. Hirsch JM, Livian G, Edward S, Noren JG. Tobacco habits among teenagers in the city of Göteborg, Sweden, and possible association with dental caries. *Swed Dent J* 1991;15:117-23.
28. Scheinin A, Mäkinen KK, Ylitalo K. Turku sugar studies. V. Final report on the effect of sucrose, fructose and xylitol diets on the caries incidence in man. *Acta Odontol Scand* 1976;34:179-216.
29. Ruxton CH, Garceau FJ, Cottrell RC. Guidelines for sugar consumption in Europe: Is a quantitative approach justified? *Eur J Clin Nutr* 1999;53:503-13.
30. Rodriguez CS, Watt RG, Sheiham A. The effects of dietary guidelines on sugar intake and dental caries in 3-year-olds attending nurseries. *Health Promot Int* 1999;14:329-35.
31. Chu YH, Tatakis DN, Wee AG. Smokeless tobacco use and periodontal health in a rural male population. *J Periodontol* 2010;81:848-54.
32. Greer RO Jr. Oral manifestations of smokeless tobacco use. *Otolaryngol Clin North Am* 2011;44:31-56, v.
33. Taybos G. Oral changes associated with tobacco use. *Am J Med Sci* 2003;326:179-82.
34. Robertson PB, Walsh MM, Greene JC. Oral effects of smokeless tobacco use by professional baseball players. *Adv Dent Res* 1997;11:307-12.
35. Tomar SL, Winn DM. Chewing tobacco use and dental caries among U.S. men. *J Am Dent Assoc* 1999;130:1601-10. Erratum in: *J Am Dent Assoc* 1999;130:1700.
36. Hsu SC, Pollack RL, Hsu AF, Going RE. Sugars present in tobacco extracts. *J Am Dent Assoc* 1980;101:915-8.
37. Talhout R, Opperhuizen A, van Amsterdam JG. Sugars as tobacco ingredient: Effects on mainstream smoke composition. *Food Chem Toxicol* 2006;44:1789-98.